



2023 Student Questions from Winters Highschool. All questions were answered by staff at Winters Healthcare.

Questions	Answer
How long does it take to find out if you are pregnant?	About six days after fertilization, the egg implants in the uterine lining. Around day 21, if conception and implantation occur during this menstrual cycle, you are pregnant.
Is it bad if cum gets in the eye?	If a person gets semen in the eyes, it will not usually cause a problem, though they may likely experience irritation or redness. People can safely flush their eyes to lower the risk of possible complications. However, in some cases, a person could develop an infection if the semen carries an STI.
Does it matter if you never actually have sexual intercourse in your life?	Sex is a personal choice. You should only have sex when you feel ready, and if that is never for you, then that is completely okay.
Why does a penis get hard when they see a hot girl?	When your brain senses a sexual urge- such as seeing a hot girl- it starts an erection by sending nerve signals to the penis. These messages tell the muscles of the corpora cavernosa (sponge-like material) to relax, allowing blood to flow in making the penis erect.
Do two condoms on top of each other work better?	Absolutely NOT. Using more than one condom at a time can cause excessive friction, and tearing of the condom can occur. This risks pregnancy and or STI transmission.
How do men cum?	“Cum” also known as semen, occurs when a penis is stimulated. This leads the penis to become erect, muscles contract, and semen is pushed out through the urethra. This is also known as ejaculation.
Do girls perform oral sex on the anus?	Anyone can perform oral sex. Whether you want to perform oral sex depends on your preferences. If you choose to perform oral sex, make sure you are using proper barrier methods to protect yourself from STIs and harmful bacteria.
Do balls get hard?	If your testicles feel hard and appear larger, it is usually because you are aroused. If you are aroused and do not get any release via orgasm, you may experience an uncomfortable, achy feeling in the



	<p>testicles known as "blue balls." Your balls do not turn blue, and you just may notice a slight blue tint because of the increased blood volume.</p>
<p>Is there a period when someone is more prone to get pregnant?</p>	<p>Yes, pregnancy most likely occurs when a person with a uterus is going through the ovulation phase of the ovarian cycle. During this time, the mature follicle is ready for fertilization. This is 12-14 days before your next period starts. Use cycle tracking.</p>
<p>What happens to a male rape victim (what should they do?)</p>	<p>ALL victims of rape should take similar steps when getting help. All individuals who are raped can get a rape kit done. Additionally, anyone who is raped should not shower to avoid washing away fluids (whether they come from someone with a penis or someone with a vagina). You should seek medical attention right away and report the sexual assault within three years. The sooner, the better!</p>
<p>What is the best position?</p>	<p>It depends on you and your partner's preference. Some positions may feel better than others. Remember to keep open communication and always get consent. Be safe and have fun!</p>
<p>Is cum good for acne?</p>	<p>There is no scientific evidence that semen is good for your skin.</p>
<p>How do you masturbate?</p>	<p>Masturbation can occur through the use of your hands, fingers, sex toys, and other objects. You can use any of the above to touch, press, rub, or massage your genitals. You may insert your fingers or a sex toy, such as a vibrator, into your vagina or anus.</p>
<p>What is the percentage of a condom working?</p>	<p>A condom for a penis is about 82% effective, and a condom for a vagina is about 79% effective. If you are not lubricating properly, friction can cause tears in the condoms, leading to ineffectiveness.</p>
<p>What does ejaculation taste like?</p>	<p>This depends on the person who is ejaculating. Diet can affect the smell of sweat, saliva, and other bodily fluids, such as breast milk, so it makes sense to assume that diet may also alter the taste of semen. However, insufficient research exists to show a strong link between diet and semen flavor. So, it depends more on the person's overall health and anatomy.</p>
<p>What does it mean to be queer?</p>	<p>To identify as queer means to identify as anything</p>




	other than straight or cisgender.
Is tearing bad when having sex for a woman? Does it require to stop?	Tearing during sex is not considered something bad. It happens to some individuals but not everyone. If tearing occurs during sex and it becomes painful, it is good to stop. Sex should be something comfortable, and communicating with your partner when this does not happen is essential. Some individuals might not feel pain during sex if there is a tear and continue having sex. Be aware of your body and stop if you feel that you should.
What is a good age to lose your virginity?	Whenever you and your partner feel ready. Do not let anyone pressure you into making this decision. If you never feel ready, that is okay too.
Does sex hurt for females?	Sex should be a pleasurable experience. Women may experience pain only once, and for others, it may be persistent. Occasional minor soreness is likely but not problematic; however, if it continues, you should seek professional advice. Don't forget to use lubrication!
What is a c-section birth, and how is it done?	A cesarean is the delivery of a newborn through a surgical incision in the abdomen and front anterior wall of the uterus. Healthcare providers use it when they believe it is safer for the baby or mother.
Why do I wake up with a morning boner?	The sex hormone testosterone plays a central role in erection. Testosterone levels fluctuate and tend to be higher early in the morning.
Is giving head good for you, or will you get sick?	There is a possibility of getting an STI through oral sex. Additionally, because all areas of the body can have bacteria on them, performing oral sex can potentially get you sick. Giving "head" is neither good nor bad; it is a personal preference whether you choose to participate in this form of sex.
How often do people get STIs unintentionally?	All the time, If someone is not practicing safe sex, not getting tested for STIs, and not communicating effectively with their partners, there is a probability that an STI can unintentionally be contracted.
How do you have sex?	Sex can mean different things to different people. Generally, sex is an activity that one, two, or more people participate in that causes arousal. Often, when people talk about sex, they are talking about



	penetrative sex. It can also involve touch, words, or both.
Why is semen clear?	Semen will normally have a clear consistency and look more white if the sperm concentration is greater. Semen consists of sperm, water, and plasma. The fluids create an environment that aids sperm in motility.
What is a micropenis?	A micropenis is a penis that is considered to be relatively small, generally about 2.5 inches or shorter. This can be classified as an infant if the penis is smaller than 0.75 inches. This is caused when the penis fails to elongate during the first trimester.
What is the average penis size?	A study of over 15,000 men found an average flaccid length of 3.6 inches and an average flaccid circumference (girth) of 3.7 inches. When erect, the penis had an average length of 5.1 inches and an average circumference of 4.5 inches.
Can you get pregnant on birth control?	Yes. Each birth control method has its percentage of effectiveness, but no method is 100% effective besides abstinence. If used correctly, some birth control methods come very close to being 100% effective at preventing pregnancy.
Can you get a person with a vagina pregnant if you ejaculate in their anus?	Ejaculation directly into the anus cannot get a person with a vagina pregnant. BUT if sperm accidentally gets on the outside of the anus and the area may be moist, sperm can travel to the vagina and cause pregnancy.
How are there more than two genders?	Gender identity is fluid. Someone can also identify as more than one gender. There are many different gender identities, including male, female, transgender, gender neutral, non-binary, agender, pangender, genderqueer, two-spirit, third gender, and all, none or a combination of these. There are many more gender identities than we've listed. Your genitals do not determine your gender identity.



<p>If you are intersex, what do you have? If you do not have a vagina or penis?</p>	<p>It depends. People who are intersex do not fully develop their genitals in the womb. So, it depends on what level the genitals developed to. Below are some examples of intersex genitalia.</p>  <p>The diagram, titled "Examples of Intersex Genital Area", shows four different anatomical views of the genital area. A legend below the diagrams identifies a blue line as "Phallosclitoris (large clitoris/small penis)" and a red line as "Labia/Scrotum".</p>
<p>Why is it so easy to throw off your pH balance?</p>	<p>It is easy to throw off the pH balance in the vagina in an acidic environment. Introducing soap or body wash with artificial scents tends to have a more basic pH; this can disrupt the good bacteria found in the vagina leading to bacterial vaginosis and other infections.</p>
<p>If you get a cold sore, do you have herpes?</p>	<p>Yes. Cold sores are a symptom of Herpes, specifically HSV-1, which is oral herpes.</p>
<p>Is it safe to have oral sex if you have had a cold sore or currently have one?</p>	<p>If you have had a cold sore in the past, but it is completely healed, and you do not currently have any open/active cold sores, then it is safe to perform oral sex on someone as it is unlikely you would pass it without an active cold sore. It is best practice to use a dental dam or a condom for oral sex to help ensure that you are protected from STIs. When cold sores are open/active, since they are herpes sores, they can still pass the virus to other places, like the genitals of a partner, if oral sex is given to someone without any protection. It is best to avoid giving oral if you have an active cold sore, but if you do, please use a condom or dental dam.</p>
<p>What is the cause of an STI?</p>	<p>An STI, or sexually transmitted disease, can be caused by bacteria or viruses.</p>
<p>How does Chlamydia happen?</p>	<p>People can get chlamydia through sexual intercourse (vaginal, oral, and anal). Chlamydia itself is a bacterial infection that can cause inflammation and</p>



	<p>irritation. Antibiotics are the type of treatment given to someone infected with Chlamydia. It is important to use the appropriate barrier methods to try and prevent getting and spreading STIs.</p>
<p>Should I have sex with a guy that hits me, but I like it?</p>	<p>You can. This would be called Sexual masochism, and it is defined as taking erotic pleasure in receiving pain. While this can include being tied up, beaten, or degraded verbally, it can also include more subtle acts, such as biting or rough sexual intercourse.</p> <p>As long as it is consensual, there are established boundaries, including a "safe word" to establish when it's too much, and your life is not at risk, then it is fine to explore your sexuality and figure out what you like.</p>
<p>I hear you can gain weight with certain birth controls. Which ones are those?</p>	<p>Everyone's body responds differently. Some birth control (typically hormonal options) may indirectly cause weight gain due to the hormones. High levels of estrogen can cause fluid retention. Increased progestin found in many forms of birth control increases appetite.</p> <p>When it comes to choosing a birth control method that is right for you, it's important to consider your personal needs.</p>
<p>Should I be scared of sex?</p>	<p>Having sex for the first time can cause anxiety. However, sex should be a pleasurable experience. Ask yourself what about sex scares you, and make a plan to address your fear. Communicate your feelings and be open about your emotions. This is especially important when working through your fears with a partner.</p>

<p>Can guys get pregnant without a surrogate?</p>	<p>Anyone who has a uterus and ovaries could become pregnant and give birth. A transgender man or non-binary (may be able to if they have a uterus)</p>
<p>Can you get an STI as a virgin?</p>	<p>The only effective method to prevent an STI is abstinence or using a barrier method such as a condom. By society, a virgin is someone who has not participated in penetrative sexual intercourse. If you are engaging in oral sex and not using a barrier, you can catch an STI.</p>
<p>Are Viagras good for you?</p>	<p>Viagra is a drug that is approved to treat erectile</p>



	<p>dysfunction. It can cause some serious side effects such as an erection that will not go away. Some more severe side effects include vision loss, hearing decrease, or irregular heartbeats. Always consult with your healthcare provider or pharmacist for other possible side effects.</p>
What does bust mean?	"Bust" is a slang term typically referring to male ejaculation/orgasm.
How long should sex last?	<p>It depends on your preference, stamina, and comfort.</p> <p>A 2008 study asked sex therapists in the United States and Canada to estimate the average duration of sex within different categories. The therapists responded that sex:</p> <ul style="list-style-type: none">• lasting under 3 minutes warrants clinical concern• lasting 3–7 minutes is “adequate”• lasting 7–13 minutes is “desirable”• lasting 10–30 minutes is “too long.” <p>However, a 2020 study assessing the time to orgasm in heterosexual women found that the average was 13.41 minutes. This suggests that the therapists polled in 2008 would consider sex that satisfies many heterosexual females to last “too long.”</p>
How many times can someone ejaculate before it becomes unsafe?	Nothing really happens when you ejaculate a lot. In fact, you can't ejaculate too much because it's self-limiting. Men are different, so they expel semen in different volumes when ejaculating. Some men don't ejaculate a lot, whereas others do. At the same time, you can't ejaculate too much in terms of a number of occasions during the day.
When you are intersex, where do you pee from?	If you are intersex, you would still urinate from your urethra, as you would if you had what is considered male or female genitalia.
What if the dad leaves?	If your partner leaves you when they find out you are pregnant, during pregnancy, or after the birth of the baby, please know that there are more people out there willing to help and support you. Parents, friends, and healthcare facilities are other sources of support for you during and after pregnancy. If your partner threatens to leave you because you are pregnant, remember you have the final say in



	<p>determining whether to remain pregnant or not. There are resources available to help during and after pregnancy, please reach out for help if you feel like you need it.</p>
Is 69'ing scary?	<p>69 refers to the slang term when two partners arrange their bodies to perform oral sex on one another at the same time. If you want to try this position with your partner, you should first have a discussion about how you would like to position yourself and what you are comfortable with.</p>
Are there internal condoms for men?	<p>Technically yes. It is like a regular condom, only larger, and it is inserted in the rectum instead of being rolled onto the penis. Inside the condom, there is a small ring that holds it inside the rectum. The internal condom is not FDA-approved.</p>
How do I respond to homophobic comments?	<p>While it is very unfortunate that some individuals choose to say homophobic or other hurtful comments please keep in mind that these people do NOT define you. Regardless of your gender identity, gender expression, who you are physically or emotionally attracted to, you are important and valued.</p> <p>If someone is saying homophobic comments that make you uncomfortable please address this with a trusted adult/teacher/parent/etc. so that conversations can be had with that individual.</p>
How to prevent STDs?	<p>To prevent STIs, you need to use a barrier method like an external or an internal condom or a dental dam. Condoms are the only contraceptive that helps prevent STIs.</p> <p>HPV is considered an STI as well. You can protect yourself from HPV by getting the three rounds of the HPV vaccine. You can also protect yourself from HIV by using PreP/PEP.</p>
Is it normal for my penis to bleed?	<p>No. Bleeding from the penis may be due to an injury, infection, or other health condition. Medication use or a long period of sexual inactivity can also cause blood to appear in sperm. It is not always a matter of concern, but some causes may need urgent medical attention.</p>
How do you safely remove the condom off of a person with a penis?	<p>Those with a penis generally lose their erection soon after ejaculating, and when that happens, the</p>



	<p>condom can slip off or spill semen. To prevent spilling, hold the base of the condom against the penis and pull both out before the penis softens. Take the condom off the penis away from your partner and throw it away.</p>
<p>What does it mean when you bleed after sex?</p>	<p>Bleeding after sex can happen if there is not enough lubrication during intercourse. Lack of lubrication can cause increased friction leading to tears in the vagina, penis, or anus. Bleeding can also happen from stretching of the hymen. This does not mean the hymen has been broken or popped, the hymen can be stretched but can also slightly tear.</p> <p>If you are constantly bleeding after sex or are experiencing pain during sex, please seek medical attention.</p>
<p>Would it be abusive/toxic for someone to keep a relationship secret if they want to come out or are in a marginalized group and would receive backlash?</p>	<p>No, it would not be abusive/toxic if someone kept their relationship a secret for the reasons mentioned. If your partner would like to keep the relationship a secret because they are not ready to come out it is important to respect your partners boundaries.</p> <p>However, you also have a say on if you are comfortable being a relationship like this.</p>
<p>Can people with a vagina/penis squirt?</p>	<p>Vagina - Squirting refers to fluid expelled from the vagina during orgasm. Not all people with vaginas squirt during orgasm, and those who do may only do it some of the time. This type of orgasm includes a <u>rapid ejection of urine, along with other fluids, from the bladder.</u></p> <p>Penis - For those with a penis, the closest equivalent to squirting is ejaculation.</p>
<p>Why does edging hurt so much?</p>	<p>Edging occurs when someone holds back their orgasm instead of releasing it. This can be painful because it is the body's natural tendency to release the tension that leads to an orgasm. Edging can also be painful in individuals with a penis because it can cause the sperm to go back up the epididymis making it uncomfortable.</p>
<p>Do penises stop growing at a certain age?</p>	<p>Penis growth begins during puberty, typically at the average age of 12 in males, and stops at the end of puberty, typically around the ages of 18-21. Penis</p>



	<p>growth, shape, and size vary greatly, and there is no “normal” size.</p>
<p>Can a penis be too big?</p>	<p>In general, most vaginas can fit all shapes and sizes of penises with proper lubrication, but it genuinely depends on your anatomy and the anatomy of your partner.</p> <p>Having a large penis can come with both physical and psychological challenges. Communicating with your partner and seeking support from medical professionals and therapists to navigate these challenges is important. Ultimately, penis size should not define or limit a person’s worth or value in society.</p> <p>It is also important to note that penis size is not the only factor that contributes to sexual pleasure and satisfaction. Communication, intimacy, and emotional connection are equally important in a fulfilling sexual experience. Therefore, it is crucial to prioritize these aspects of a relationship rather than solely focusing on physical attributes.</p>
<p>What is head?</p>	<p>Head and/or Giving Head is a slang term to reference oral sex.</p>
<p>Why can some people not get pregnant?</p>	<p>There are a lot of factors that contribute to if someone can get pregnant. Some common reasons someone can not get pregnant are -</p> <ul style="list-style-type: none">You might not have been trying long enough.You might not be ovulating.Your male partner may be experiencing infertility.You may be experiencing age-related infertility.You may have blocked fallopian tubes.You may have endometriosis.You may have underlying medical problems.You may be on medications that impact fertility.You may have lifestyle factors that impact fertility.You could have unexplained fertility.
<p>Is it normal for your period to vary based on month?</p>	<p>Yes, periods vary from individual to individual. While some can have an almost identical flow each month this is not the case for most individuals with a uterus.</p> <p>Additionally, the menstrual cycle is dependent on the release of hormones, if someone is experiencing stress (emotional, physical, nutritional, etc.) this can have a direct effect on the release of hormones in the</p>



	<p>body. If your period tends to be very abnormal or extremely painful please visit your medical provider. Abnormal periods can also make it difficult to track ovulation cycles leading to higher risk of unplanned pregnancy.</p>
<p>How probable is it for a condom to break?</p>	<p>The odds of a condom breaking are <u>small if you're using them correctly</u>, but it can definitely happen. While statistics vary, if used correctly it is about a 1-12% chance it could break. Here are some tips on how to help ensure your condom doesn't break-</p> <ul style="list-style-type: none">• Store condoms in a cool, dry place (so not in your wallet or very tight pockets for weeks or months).• Check the expiration date to make sure the condoms you're using are still good.• Use water or silicon-based lube. Lube reduces the friction that can lead to breakage.• Make sure you put the condom on before you start having sex, and keep wearing it until you've finished and pulled out.
<p>Does using a tampon/inserting a cup in for a period make you "not a virgin"?</p>	<p>NO. The concept of being a virgin is a societal-based concept/ideal. Your anatomy does not change after sex. The hymen is stretchy and elastic, like a hair scrunchy. It is meant to be able to expand and retract. The Hymen is not a seal or barrier that needs to be broken as it is thought when someone loses their virginity / "gets their cherry popped." There is nothing to "pop". So, to insert something like a tampon into your vagina does not take your "virginity."</p>
<p>When you use a condom, do you still use a Plan B?</p>	<p>If you use an Internal or an external condom. It doesn't break and is used correctly; you should not need to take Plan B or any other emergency contraceptive. Keep in mind that it is ideal to use a condom even if you are on another form of birth control that doesn't prevent STIs like the pill, patch, NuvaRing, etc.</p> <p>If the condom breaks or is not removed correctly and semen does enter the vaginal canal, then you SHOULD take an emergency contraceptive as soon as possible.</p>
<p>Is a clitoris basically a penis?</p>	<p>Sort of. It doesn't have a central role in reproduction like the penis or vagina — it's basically just there to</p>



	<p>make you feel good! The clitoris has thousands of nerve endings in it, which makes it super sensitive. It can be a little tricky to find your clitoris, since it's at the top of your vulva, tucked inside your inner labia, underneath its own protective hood. For some people the clitoris is easily visible and accessible, and for others it's hidden under the clitoral hood – everyone's vulva is unique. The clitoris can vary in size (it can be about as small as a pea or as big as a thumb), but only the tip of it comes outside the body where you can see it. When you're aroused (turned on), your clitoris becomes stiff and swollen — <u>just like a penis</u>. If you're having trouble finding it, things like looking at and touching your own vulva can help you figure out how your vulva is built, and what feels good. Fun fact about the clitoris: while you can only see the very tip of it, it actually has a spongy shaft that stretches up to 5 inches inside your body!</p>
<p>What are the common side effects of taking birth control?</p>	<p>It depends on which birth control, as well as how your body reacts to that specific birth control.</p> <p>Some common side effects are</p> <ul style="list-style-type: none">• Minor weight gain, typically related to increased appetite or fluid retention• Decreased libido• Headaches• Nausea• Changes in your period.
<p>After having your period for two years, is it normal to still miss 1-2 months?</p>	<p>If your periods are consistently irregular, it is recommended you see your doctor or gynecologist. Your menstrual cycle may not always be predictable, and that may be okay. However, look out for periods that occur fewer than 21 days or more than 35 days apart. Missing three or more periods in a row may be considered irregular. Hormone levels, certain health conditions, and medications can contribute to an irregular period.</p>
<p>Can you get pregnant only from penetration?</p>	<p>No, pregnancy can occur if any sperm enters the vaginal canal and fertilizes an ovum, even if it did not occur through penetration.</p> <p>Because sperm are mobile and thrive in moist environments, pregnancy can occur if anal sex is performed and sperm makes it to the vaginal canal. If sperm is ejaculated on top of an individual with a</p>



	<p>uterus close to but not inside the vagina, there is still a possibility that it can make it to the vaginal canal if the area is moist.</p>
<p>Can you get pregnant easily?</p>	<p>While getting pregnant can be easy for some people it can be more difficult for others. It depends on the individual, their overall health, and family history.</p> <p>If you have considered the cost of raising a child, your living situation, your financial stability, etc. and are prepared to have a kid it is important to remember that there are many different methods of having a child. Even if you are not able to conceive one yourself. Ex's</p> <ul style="list-style-type: none">● IVF (In Vitro fertilization)● Surrogacy● Adoption● Artificial Insemination● And many others!
<p>How does chlamydia happen?</p>	<p>You can get chlamydia by having vaginal, anal, or oral sex with someone who has chlamydia. Also, you can still get chlamydia even if your sex partner does not ejaculate (cum). A pregnant person with chlamydia can give the infection to their baby during childbirth.</p>
<p>Can you get supplies for free if you don't have sufficient funds?</p>	<p>Yes! Absolutely.</p> <p>Winters Healthcare has a Family Pact program that allows you to access free and confidential (<i>meaning your parents would not know</i>) services.</p> <p>Family Pact Covers -</p> <p>Various birth control methods, including condoms, pills, long-acting reversible contraceptives, emergency contraception, and sterilization</p> <ul style="list-style-type: none">● Family planning counseling and education● Sexually transmitted disease (STD) testing & treatment● HIV testing● Cervical cancer screening● Limited fertility services



Can a condom break during sex?	<p>Yes.</p> <p>There are various reasons a condom can break. If there isn't enough lubrication, if the condom is expired, or if the condom has been stored improperly, these are just some of the reasons why it may break. Always check after to make sure the condom doesn't break.</p>
Should I be scared of sex?	<p>It is expected to be nervous about having sex, especially if it is the first time. Sex should be a pleasurable and fun experience. It should never be forced or scary.</p>
What is the cause of an STI?	<p>STI's are mainly spread through unprotected sexual contact in blood, semen, or vaginal and other bodily fluids. They are caused by bacteria, viruses, or parasites.</p>
How do you safely remove a condom from a penis?	<p>Guys generally lose their erection soon after ejaculating, and when that happens it's possible for the condom to slip off or spill semen. To prevent spilling, hold the base of the condom against the penis and pull both out before the penis softens. Take the condom off the penis away from your partner and throw it away.</p>
What should you do if a condom breaks in you to stop pregnancy?	<p>If a condom breaks and you want to avoid getting pregnant, take an emergency contraceptive right away. This can be either Plan B (take within 72 hours) or Ella (take within 5 days). Remember, <u>the sooner you take it, the better it works</u>. The pharmacy at Winters Healthcare sells Plan B for \$10.</p> <p>Please keep in mind, Plan B is not a primary form of birth control.</p> <p>If a condom breaks it is also super important to get tested for STI's.</p>
Is it normal for my penis to bleed?	<p>No. The only normal discharge should be urination or semen/ejaculation for those with a penis. If you have blood or other abnormal discharge, please seek medical attention.</p>
What is the most accessible form of birth control without parental consent?	<p>You do not need parental consent for most forms of birth control. Winters Healthcare has a Family Pact program that allows you to access free and confidential (<i>meaning your parents would not know</i>) services.</p>



	<p>Family Pact Covers - Various birth control methods, including condoms, pills, long-acting reversible contraceptives, emergency contraception, and sterilization.</p> <p>Call Winters Healthcare and set up an appointment and come talk with a provider to pick which method is best for you and your lifestyle. No parental consent needed!</p>
Should I have sex with a gang banger?	<p>You can have sex with whomever you choose. Sex must always be consensual. Communicate, and be aware of your sexual partner's lifestyle, and always use an internal or external condom to avoid an STI or unwanted pregnancy.</p>
What if the condom is too big or too small?	<p>There are different sizes of condoms available. The standard-sized condoms are about 7.8 inches. Larger condoms are about 8.1 inches. Ensure that the condom you are using fits comfortably to lessen the chance of the condom breaking or slipping off. It should be a bit snug, but not too tight, cutting off circulation.</p>
Do you get your period with birth control?	<p>This depends on the type of birth control. In general, women who use the IUD, patch, pill, or ring will have menstrual periods every month. Women who take the shot may not have periods often or at all while they use the method. Some with a uterus choose not to have a period on the pill, patch, or ring. They do this by skipping the "break" (like the sugar pills in the pill pack) in birth control that allows you to menstruate, and that is totally safe and fine!</p> <p>Make sure to tell your provider you would like to skip your period so that they can prescribe you enough of that birth control to do so.</p>
Can a condom get stuck inside of you?	<p>The vagina has a stopping point which is the cervix, so condoms cannot get lost inside the pelvic cavity, but they can get lodged towards the top of it the cavity (by the cervix) and go unnoticed.</p> <p>Condoms may slip off during sex if they are not the right size or if not put on properly. If a condom does get stuck inside of the vagina, you will need to get it out so it does not cause problems such as an STI or unplanned pregnancy. If you can't remove the</p>



	<p>condom completely, call your doctor or a health center right away.</p>
<p>Can you sue the condom company if it rips?</p>	<p>You can, but you would have to prove you did everything correctly and it still broke. Prophylactic Liability. The condoms will likely have a warning or a disclaimer on the consumer package to protect the company against legal action. So be aware that the product may have functioned as advertised, even if it broke during the use. However, if a condom is poorly designed or if it malfunctions, the condom manufacturer could be held responsible under product liability law. Condom manufacturers have a duty to ensure their products are safe, and that they function as advertised. Normally it's hard to find the evidence to prove the condom's quality meet problem or the misuse of it. Normally condom's quality is 100% assured, most of the failure caused by the wrong use.</p> <p>Carefully read the instructions before use a condom, this will make sure the max assurance to protect you.</p>
<p>What flavor of condoms are there? Are the colors and flavors matching? Example: grapes are purple.</p>	<p>Flavored condoms are designed to be used for oral sex and help mask the taste of latex. There are a wide variety of flavors such as vanilla, chocolate, strawberry, etc. The color of the packaging does not necessarily indicate the flavor however some brands do have matching flavors to colors.</p>
<p>What are the side effects of Nexplanon?</p>	<p>The most common implant side effect is spotting (light bleeding) discharge. This is common in the first 6-12 months. Although rare, implants can cause long-term spotting or heavier periods. For most, the implant makes periods lighter or even makes it stop (this is normal). Other possible side effects include Headaches, breast pain, nausea, weight gain, ovarian cysts, pain or bruising at the site of implant or infection.</p>
<p>Is it illegal to have sex with an 18-year-old if you are 14 years old?</p>	<p>Having sex with a minor is considered statutory rape. Even if you give consent as a 14-year-old.</p>
<p>What does anal mean?</p>	<p>Anal refers to the act of penetration of the anus.</p>
<p>How can you tell when a condom rips?</p>	<p>It is often hard to know during intercourse if the condom has ripped. That is why it is essential to check before you use a condom and after intercourse with a condom if the condom is broken or not.</p>



Can I get a pink condom?	Yes. There are a wide variety of condoms, which include colors, flavors, thickness, and size.
Does the pull-out method work?	The pull-out method/withdrawal works about 78% of the time overall. But the odds of pregnancy are always higher during the 5 days leading up to, and during ovulation — these are called fertile days. If no semen gets on your vulva or in your vagina, pregnancy can't happen — whether or not you're ovulating.
How do you use a vibrator? Does it hurt?	<p>Using a vibrator comes down to your sexual preferences and what makes you feel comfortable. Find a setting you think is fun and use it on different parts of your body to see how it feels. Remember, there's no 'right' way to pleasure yourself.</p> <p>Vibrators are typically used by placing the device on sensitive erogenous zones, which are places on your body that increase sexual pleasure. They can be used externally on the penis, vulva, clitoris and anus, or can be used internally in the anus or vagina. It should not hurt to use a vibrator, ensure you are using lubricant and try to relax. If it does hurt, that may not be the vibrator for you!</p> <p>According to a study by the U.S. Library of Medicine, vibrations increase blood flow and improve both pelvic muscle tone and relaxation, and they're more likely to help you orgasm.</p> <p>Overall, make sure you read the directions, check that the vibrator is cleaned properly, and use a PH based lubricant that is safe for use with toys.</p>
Does putting a tampon in feel the same as vaginal penetration?	Not typically, no. There are different components and emotions being felt when someone is performing sex compared to when someone is putting in a tampon. Using tampons is not associated in any way to the loss of "virginity".



<p>Is it true your hips get bigger with sex?</p>	<p>No, this is not true. Sex won't change the way your body looks, because there's zero connection between body growth and sexual activity. It is true that some young women start having sex around the time those changes take place. So they may think that having sex causes the changes, but it's just coincidence. However, pregnancy — which can result from sex — does cause changes in a woman's body.</p>
<p>If a woman sits on top of a penis, can it break?</p>	<p>A penis does not break like the bones in your body. The spongy tissue within the penis surrounding the urethra is known as the corpus spongiosum. When a penis is erect, a penile fracture can occur but it is rare.</p>
<p>Is it bad to have sex more than one time a day?</p>	<p>As long as no one is being pressured to have sex more than they want to, there's nothing wrong with having sex as often as feels right for you and your partner. Just make sure you don't forget to take care of yourself, your partner, and to listen to your body.</p>